STATE UNIVERSITY OF NEW YORK COLLEGE OF OPTOMETRY ®

BACKGROUND

Despite recent advancements in laser refractive procedures, including the introduction of small-incision lenticule extraction (SMILE), some postrefractive surgery patients still experience myopic regression over time. While the mechanism of regression following refractive treatment is not fully understood, the process is clearly multifactorial, involving the corneal epithelium and stroma.¹

Amongst numerous treatment modalities for myopia, orthokeratology (orthoK) is capable of satisfying patients' visual needs with low risk, thereby benefiting the same group of highly motivated potential refractive surgery candidates. Orthok is a non-surgical, reversible treatment for myopic refractive error which flattens the cornea temporarily using a specially designed reverse-geometry contact lens. Although orthokeratology is primarily fit on adolescents to deter myopia progression and to provide refractive correction, it is proven to correct myopia in both adult and pediatric patient populations. For post-refractive surgery patients who are interested in continued freedom from daytime glasses or contact lenses, orthokeratology is a reasonable alternative to enhancement surgery. Successful management of post-refractive surgery myopic regression with orthokeratology has been reported.² This case study explores the visual performance of a post-SMILE patient who was successfully treated with orthokeratology.

CASE

A 24-year-old Asian female presented to the University Eye Center complaining of distance blur OS five years s/p SMILE surgery OU. Pre-SMILE, she successfully wore Paragon CRT® lenses:

	OD	OS
Refraction	-5.25 -0.50 x 180	-5.25 -1.75 x
Lens Parameters	9.20 - 525 - 32	9.20 – 550 -
BCVA	20/20	20/20

Table 1: Patient's pre-SMILE refraction and orthoK parameters

The patient discontinued contact lens wear, as directed, for six months before proceeding with SMILE OU.





Figure 1: Pre-SMILE tangential topography maps

Post-SMILE Regression

Despite a successful outcome post-operatively (20/20 vision OD, OS), patient's myopia had progressed in the left eye alone over the last two years. Uncorrected visual acuity (UCVA) was 20/20 OD, 20/40 OS and manifest refraction was plano OD and -1.50 sph OS at the time of initial consultation.



Figure 2: Post-SMILE, pre-orthoK topography maps





Enhance your SMILE with Orthokeratology: A Case Report on the Use of **Orthokeratology to Treat Myopia Regression After SMILE Surgery**

Shuyi (Suzy) Chen*, Elton Wong, Sharon P. Keh OD FAAO



Patient was motivated to pursue orthokeratology to address her declining distance acuity OS, as she had worn Paragon CRT® lenses prior to SMILE surgery.

Lens 1 (Initial Lens)

Based on patient's refraction and topography, the following Paragon CRT Dual Axis ® lens was ordered empirically with consultation.



Figure 3: Lens 1, Day 1 topography and slit lamp photo OS

Day 1 follow-up of Lens 1 revealed superior temporal decentration, excessive pooling in the RZD and a significant over-refraction. The lens required the following modifications: reducing the RZD and flattening the BCR to decrease the sagittal depth, achieve better centration, and address the lack of treatment.

Lens 2



Figure 4: Lens 2, Day 1 topography and slit lamp photo OS

While the second lens improved vision significantly, the treatment zone still exhibited superior temporal decentration. Therefore, BC was further flattened and lens diameter increased in an attempt to provide better uncorrected VA and improve lens centration.

Lens 3 (Final Lens)



Figure 5: Lens 3, Day 1 topography and slit lamp photo OS

An ideal fluorescein pattern with an adequate and centered treatment zone was achieved with the final lens. Patient was pleased with the level of uncorrected vision and enhanced stability of refractive state throughout the day.

LZA	Diameter	
31	11.0	
and a start		
		Day 1
	UCVA	20/40
	Refraction	-1.00 sph
	Table 2: Lens 1, Day 1	refraction OS

LZA	Diamete	r	
31	11.0		
		Day 1	Week 1
•	UCVA	20/30	20/25+2
	Refraction	-0.75 sph	pl -0.75 x 90
	Table 3: Lens 2, Day 1 and Week 1 refraction OS		

Diameter		
11.5		
	Day 1	Week 1
UCVA	Day 1 20/25 ⁻²	Week 1 20/20

DISCUSSION

Fitting post-refractive surgical patients with orthokeratology can be challenging because of their iatrogenic, oblate corneal shape. In this case, proper centration, fluorescein pattern, and desired myopia reduction were achieved with the customization of a Paragon CRT Dual Axis[®] lens. This case study also investigated the effect of orthoK treatment on additional factors which may affect visual performance. At each follow up visit of the final lens, patient's contrast sensitivity was assessed with Pelli-Robson contrast sensitivity chart and corneal aberrations were measured on the OPD-Scan III Wavefront Aberrometer.

Corneal Spherical Ab Corneal Higher Order Log Contrast Sensi

Table 5: Patient's contrast sensitivity and corneal aberrations

After 1 day and 1 week of final lens wear, an increase in both corneal spherical aberration and higher order aberration were observed, as well as a slight decrease in contrast sensitivity. These progressions are supported by previous research on orthokeratology, which has been shown to induce corneal aberrations causing a decrease in contrast sensitivity function.³ Changes in these parameters, particularly the increased aberrations and reduced contrast sensitivity, can affect overall visual performance. It is important to note that patients may experience symptoms not only from orthok treatment but from previous refractive surgery. Pre-SMILE corneal aberrations and contrast sensitivity were not acquired for this case but would have been interesting data for comparison's sake.

Fortunately, our patient maintained superb subjective vision throughout the day. Close follow-up is indicated in post-refractive surgery patients to address possible visual complaints and to monitor the status of their post-surgical cornea. More research is warranted on long-term visual performance of post-refractive surgery patients treated with orthoK, and the role of corneal aberrations and contrast sensitivity in success.

CONCLUSION

It is possible to successfully fit post-SMILE patients with orthokeratology to correct myopia secondary to regression, despite the relatively flat central corneal curvature with respect to the steeper mid-peripheral cornea. This case suggests that overnight orthokeratology is a safe and effective alternative to enhancement surgery.

REFERENCES

<u>0914</u>.

ACKNOWLEDGEMENTS

Special thanks to Paragon Vision Sciences, Inc. for generously providing the lenses for this case study and, in particular, Kelly Voltz, OD, FAAO, for her invaluable advice and guidance throughout the case.

	Doct SNALLE Dro Orthok	Lens 3 (Final lens)		
	Post-Siville, Pre-Orthok	Days 1	Week 1	
erration	0.020	0.037	0.056	
berration	0.109	0.139	0.157	
itivity	1.68	1.64	1.60	

L. Moshirfar, Majid et al. "Mechanisms of Optical Regression Following Corneal Laser Refractive Surgery: Epithelial and Stromal Responses." Medical hypothesis, discovery & innovation ophthalmology journal vol. 7,1 (2018): 1-9. 2. Park, Young Min et al. "Effect of Orthokeratology in Patients with Myopic Regression after Refractive Surgery." Contact Lens and Anterior Eye 39, no. 2 (2016): 167–71. https://doi.org/10.1016/j.clae.2015.10.003.

^{3.} Hiraoka, Takahiro et al. "Contrast Sensitivity Function and Ocular Higher-Order Aberrations Following Overnight Orthokeratology." Investigative Ophthalmology & Visual Science 48, no. 2 (2007): 550. https://doi.org/10.1167/iovs.06-